

# Welcome to Physics

#### Chapter 26: Properties of Light

www.MrLiddell.com

# **Convert the following into Essential Questions:**

- Electromagnetic Waves
- The Electromagnetic Spectrum
- Transparent Materials
- Opaque Materials
- Seeing Light—The Eye

#### **Electromagnetic Waves**


- Light is the only thing we can see.
  - Originates from the accelerated motion of electrons
  - Electromagnetic phenomenon



#### **Electromagnetic Waves**

#### Electromagnetic wave

 Made up of vibrating <u>electric and magnetic</u> fields



# Electromagnetic Waves CHECK YOUR NEIGHBOR


If an electron vibrates up and down 1000 times each second, it generates an electromagnetic wave with a

- A. period of 1000 s.
- B. speed of 1000 m/s.
- C. wavelength of 1000 m.
- D. None of the above.

#### **Electromagnetic Spectrum**

#### Electromagnetic spectrum

- Classification of electromagnetic waves according to frequency
  - Lowest frequency of light we can see appears red.
  - Highest frequency of light we can see appears violet.
  - Higher frequency of light is ultraviolet—more energetic and causes sunburns.
  - Beyond are X-ray and gamma ray.



© 2015 Pearson Education, Inc.

# Electromagnetic Spectrum CHECK YOUR NEIGHBOR

The electromagnetic spectrum spans waves ranging from lowest to highest frequencies. The smallest portion of the electromagnetic spectrum is that of


- A. radio waves.
- B. microwaves.
- C. visible light.
- D. gamma rays.

# Electromagnetic Spectrum CHECK YOUR NEIGHBOR


Which of these is fundamentally different from the others?

- A. Sound waves
- B. Light waves
- C. Radio waves
- D. X-rays

- Light is transmitted similarly to sound.
  - Both are vibrations due to a vibrating source.




 How light penetrates transparent material such as glass:



- How light penetrates transparent material such as glass (continued)
  - Electrons or molecules in the glass are forced into vibration.
  - Energy is momentarily absorbed and vibrates the electrons in the glass.
  - This vibrating electron either emits a photon (a corpsucle of light) or transfers the energy as heat.
- Time delay between absorption and re-emission of energy of vibrating electrons results in a lower average speed of light through a transparent material.

- In glass, infrared waves, with frequencies lower than those of visible light, cause not only the electrons but entire atoms or molecules to vibrate, increasing the temperature of the structure.
- So we see that glass is transparent to visible light, but not to ultraviolet and infrared light.



- Average speed of light through different materials
  - vacuum—c (300,000,000 m/s)
  - atmosphere—slightly less than c (but rounded off to c)
  - water-0.75 c
  - glass—0.67 c, depending on material
  - diamond—0.41 c

# Transparent Materials CHECK YOUR NEIGHBOR

Strictly speaking, the photons of light incident on glass are

- A. also the ones that travel through and exit the other side.
- B. not the ones that travel through and exit the other side.
- C. absorbed and transformed to thermal energy.
- D. diffracted.

# Transparent Materials CHECK YOUR NEIGHBOR

Compared with the frequency of illuminating light on a sheet of transparent plastic, the frequency of light that is transmitted

- A. is slightly less.
- B. is the same.
- C. is slightly higher.
- D. depends on the type of plastic.

# Transparent Materials CHECK YOUR NEIGHBOR

The average speed of light is less in

- A. air before entering glass.
- B. glass.
- C. air after emerging from glass.
- D. None of the above.

 Most things around us are opaque—they absorb light without re-emitting it.

- Books, desks, chairs, and people are opaque.

 Vibrations given by light to their atoms and molecules are turned into random kinetic energy—into internal energy.

These materials become slightly warmer.

- Metals
  - Light shining on metal forces free electrons in the metal into vibrations that emit their own light as reflection.




- Light incident on
  - dry surfaces bounces directly to your eye.
  - wet surfaces bounces inside the transparent wet region, absorbing energy with each bounce, and reaches your eye darker than from a dry surface.

#### Shadows


- A thin beam of light is often called a *ray*.
- When we stand in the sunlight, some of the light is stopped while other rays continue in a straight-line path.
- We cast a shadow—a region where light rays do not reach.

- Either a large, far-away light source or a small, nearby light source will produce a sharp shadow.
- A large, nearby light source produces a somewhat blurry shadow.




© 2015 Pearson Education, Inc.


- Light is the only thing we see with the most remarkable optical instrument known—the eye.
- As light enters the eye, it moves through the transparent cover called the *cornea*, which does about 70% of the necessary bending of the light before it passes through an opening in the *iris* (colored part of the eye).




- The opening is called the *pupil*.
- The light then reaches the *crystalline lens,* which finetunes the focusing of light that passes through a gelatinous fluid called *vitreous humor.*
- Light then passes to the *retina,* which covers the back two-thirds of the eye and is responsible for the wide field of vision that we experience.



- For clear vision, light must focus directly on the retina.
- The retina is not uniform.
  - In the middle is the macula, and a small depression.
  - in the center is the *fovea*, the region of most distinct vision.
  - Behind the retina is the optic nerve, which transmits signals from the photoreceptor cells to the brain.
  - There is also a spot in the retina where optic nerves are connected; this is the blind spot.




- The retina is composed of tiny antennae that resonate to the incoming light.
- Rods handle vision in low light.
  - They predominate toward the periphery of the retina.
- **Cones** handle color vision and detail.
  - They are denser toward the fovea.
  - There are three types of cones, stimulated by low, intermediate and high frequencies of light.



Although our vision is poor from the corner of our eye, we are sensitive to anything moving there.




- The brightest light that the human eye can perceive without damage is some 500 million times brighter than the dimmest light that can be perceived.
- Lateral inhibition: We don't perceive the actual differences in brightness.
  The brightest places in our visual field are prevented from outshining the rest.






- The brightest light that the human eye can perceive without damage is some 500 million times brighter than the dimmest light that can be perceived.
- Lateral inhibition: We don't perceive the actual differences in brightness.
  The brightest places in our visual field are prevented from outshining the rest.











Are the dashes on the right really shorter?

© 2015 Pearson Education, Inc.



Can you count the black dots?




© 2015 Pearson Education, Inc.




Is the hat taller than the brim is wide?



What does this sign read?



Are the rows of tiles really crooked?



Are the vertical lines parallel?